

Computational Geometry Column 44

Joseph O'Rourke*

Abstract

The open problem of whether or not every pair of equal-area polygons has a hinged dissection is discussed.

A *dissection* of one polygon A to another B is a partition of A into a finite number of pieces that may be reassembled to form B . A and B are then said to be *equidecomposable*. A staple of puzzle collections for centuries, the field has been given new life through Frederickson's 1997 monograph [Fre97]. He quickly followed that with a second book [Fre02] on *hinged dissections*, the most famous of which is the elegant 1902 Dudeney-McElroy hinged dissection between a square and an equilateral triangle illustrated in Fig. 1.




Figure 1: A four-piece hinged dissection between a square and an equilateral triangle.

It has been known since the early 19th century that any two polygons of equal area are equidecomposable.¹ One method [Fre97, p. 221] triangulates polygon A , dissects each triangle to a rectangle, converts each rectangle to one of fixed length, and stacks all the rectangles. Applying the same process to B and overlaying the stacked rectangles yields a mutual dissection. No comparable method is known for hinged dissections:

*Dept. of Computer Science, Smith College, Northampton, MA 01063, USA. orourke@cs.smith.edu. Supported by NSF Distinguished Teaching Scholar Grant DUE-0123154.

¹It has been known since the early 20th century that not all equal-volume polyhedra are equidecomposable.

QUESTION: Does every pair of equal-area polygons have a hinged dissection?

The usual interpretation of this problem ignores possible intersections between the pieces as they hinge, following what Frederickson calls the “wobbly-hinged” model. Removing this freedom is perhaps not difficult, but seems not to be settled in the literature.

Here I report on two advances on this problem. The first is a proof that any two *polyominoes* of the same area have a hinged dissection [DDE⁺03]. A polyomino is a polygon formed by joining unit squares at their edges [Gol65, Kla97]. One composed of n squares is called an n -omino. The authors establish a type of universal dissection by proving that a cycle of $2n$ right isosceles triangles, hinged at their base vertices, can fold to any n -omino. See Fig. 2. Although the cycle may seem a special case, they prove that any hinged graph structure may be converted into another dissection hinged in a cycle. The polyomino result




Figure 2: Eight right isosceles triangles hinging to a tetramino.

generalizes to hinged dissections of all edge-to-corresponding-edge gluings of congruent copies of any polygon. This leads, among other curiosities, to a 128-piece hinged chain of isosceles right triangles that folds to shapes representing all the letters of the alphabet and the ten digits [DD03].

The second advance is a proof by Eppstein [Epp01] that any asymmetric polygon has a hinged dissection to its mirror image. His dissection partitions the polygon into a chain of *kites*, quadrilaterals with reflection symmetry across a diagonal. With this partition in hand, it is easy to rotate the kites so all their symmetry diagonals align. The mirror image of the unfolding process then folds the shape to its mirror image. This procedure is illustrated for a simple example in Fig. 3. The kite partition relies on circle packing [BE00], and must be supplemented by a refined partition, and threading around the boundary of a spanning tree to obtain a hinged chain. This work leads to a possible route toward solving the general problem for hinged dissections, for it reduces it to $O(n)$ equal-area triangle hinge-dissection problems with particular constraints




Figure 3: A dissection into a chain of kites, hinging to the mirror image. Here the kites are all squares.

on the hinges.

References

- [BE00] Marshall W. Bern and David Eppstein. Quadrilateral meshing by circle packing. *Int. J. Computational Geometry & Applications*, 10(4):347–360, August 2000.
- [DD03] Erik D. Demaine and Martin L. Demaine. Hinged dissection of the alphabet. *Journal of Recreational Mathematics*, 31(3):204–207, 2003.
- [DDE⁺03] Erik D. Demaine, Martin L. Demaine, David Eppstein, Greg N. Frederickson, and Erich Friedman. Hinged dissection of polyominoes and polyforms. *Computational Geometry: Theory and Applications*, 2003. To appear. arXiv:cs.CG/9907018. Revised version of paper in *Proc. 11th Canad. Conf. Comput. Geom.*, 1999, 15–18.
- [Epp01] David Eppstein. Hinged kite mirror dissection. ACM Computing Research Repository, June 2001. arXiv:cs.CG/0106032.
- [Fre97] Greg Frederickson. *Dissections: Plane and Fancy*. Cambridge University Press, 1997.
- [Fre02] Greg Frederickson. *Hinged Dissections: Swinging & Twisting*. Cambridge University Press, 2002.
- [Gol65] Solomon W. Golomb. *Polyominoes*. Charles Scribner’s Sons, New York, 1965.
- [Kla97] David A. Klarner. Polyominoes. In Jacob E. Goodman and Joseph O’Rourke, editors, *Handbook of Discrete and Computational Geometry*, chapter 12, pages 225–242. CRC Press LLC, Boca Raton, FL, 1997.